

Final Progress Report

United States Army Research Office, Terrestrial Sciences

Grant Number -- DAAH04-96-1-0036

Principal Investigator -- Dr. Paul R. Bierman

Statement of Problem

Currently, evaluating long-term rates of land surface erosion is a very difficult and uncertain task. Knowing such rates is imperative if land management decisions are to be made responsibly. Only if land managers can compare human-induced rates of surface change, with natural or background rates, can management decision be made cost-efficiently and with the best balance between preservation and use of the land.

In order to further this goal, we determined the background or natural rate at which soil and rock surfaces are eroding at three different sites: the Yuma Proving Ground (Yuma, AZ), the southern Mojave desert (near General Patton's former camp at Iron Mountain), and at Nahal Yael (an intensively studied, hyper-arid basin in the southern Negev of Israel). Such information is useful to land managers who need a context in which to evaluate responsibly the impact of Army mission activities.

Summary of Most Important Results

This funding supported part-time salary, field, and analytic costs for seven students, a faculty member, and a technician to collect and analyze samples from two sites: Yuma Proving Ground, California and the Nahal Yael Research Watershed, Israel. In addition, salary support for Bierman and Clapp was used for the analysis of other related data sets including sediment samples collected from similar landscapes and modelling of complex exposure histories. The results of all of this research are reported in detail by the papers and abstracts listed in the next section.

In summary, our most important findings are:

- At Yuma Proving Ground, we determined that most sediment in upland drainages is derived from hillslopes and that rocky outcroppings are the most stable landscape features. This suggests that while rocky outcrops may be the landscape features most resistant to erosion they will also be the slowest features to recover from human impact.
- On the main stem of Yuma Wash, we determined that much of the sediment transported by the Wash is actually derived not from erosion of the highlands but from reworking of the older valley filling alluvium. This finding is important in terms of land management if the need arises to limit the sediment discharge from Yuma Wash. Limiting such discharge will require erosion control not only in the highlands but also in the valley.

REPORT DOCUMENTATION PAGEForm Approved
OMB NO. 0704-0188

Public Reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188.) Washington, DC 20503.

1. AGENCY USE ONLY (Leave Blank)		2. REPORT DATE June 2000		3. REPORT TYPE AND DATES COVERED Final Report	
4. TITLE AND SUBTITLE Estimating Long Term Rates of Erosion and Sediment Generation Using Cosmogenic Isotope Abundances in Rock, Soil and Sediment				5. FUNDING NUMBERS DAAH04-96-1-0036	
6. AUTHOR(S) Paul R. Bierman					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Vermont Burlington, VT 05405				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U. S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211				10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 34572.2-EV	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.					
12 a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.				12 b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) Currently, evaluating long-term rates of land surface erosion is a very difficult and uncertain task. Knowing such rates is imperative if land management decisions are to be made responsibly. In order to further this goal, we determined the background or natural rate at which soil and rock surfaces are eroding at two sites: the Yuma Proving Ground (Yuma, AZ) and at Nahal Yael (an intensively studied, hyper-arid basin in the southern Negev of Israel). At Yuma Proving Ground, we determined that most sediment in upland drainages is derived from hillslopes and that rocky outcroppings are the most stable landscape features. On the main stem of Yuma Wash, we determined that much of the sediment transported by the Wash is actually derived not from erosion of the highlands but from reworking of the older valley filling alluvium. At Nahal Yael watershed, we that long-term sediment generation rates are substantially less than short term sediment yields as measured by trapping the last 30 years of sediment transported out of the basin behind a retention dam. These findings suggest that short-term measurements of sediment yield may be overestimates.					
14. SUBJECT TERMS erosion, sedimentation				15. NUMBER OF PAGES 3	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED	18. SECURITY CLASSIFICATION ON THIS PAGE UNCLASSIFIED	19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED	20. LIMITATION OF ABSTRACT UL		

NSN 7540-01-280-5500

Standard Form 298 (Rev.2-89)
Prescribed by ANSI Std. Z39-18
298-102

DTIC QUALITY INSPECTED 4

- At Nahal Yael watershed, we found a similar pattern of stable bedrock and more rapidly eroding hillslopes. We also found that long-term sediment generation rates are substantially less than short term sediment yields as measured by trapping the last 30 years of sediment transported out of the basin behind a retention dam. These findings suggest that short-term measurements of sediment yield may be overestimates. Such a finding has important implications for the design of long-lasting engineering works such as dams and sedimentation control structures.

List of Publications (supported in whole or in part by ARO funding of salary and/or analyses)

Refereed papers

- Bierman, P., Clapp, E.M., Nichols, K.K., Gillespie, A.R., Caffee, M. (in prep.) Using cosmogenic nuclide measurements in sediments to understand background rates of erosion and sediment transport, *in* Landscape Erosion and Evolution Modelling, Harmon, R. S. and Doe, W. M., eds., Kluwer.
- Clapp, E. M., Bierman, P.R., Schick, A. P. Lekach, J., Enzel, Y., and Caffee, M., (in review), Differing rates of sediment production and sediment yield, **Geology**.
- Clapp, E., Bierman, P.R., and Caffee, M. (in review) Using ^{10}Be and ^{26}Al to determine sediment generation rates and identify sediment source in an arid region drainage basin, **Geomorphology**.
- Nichols, K.K., Bierman, P. R., and Caffee, M., (in review) Understanding desert Piedmonts using in situ produced ^{10}Be and ^{26}Al , **Geomorphology**.
- Clapp, E., Bierman, P.R., Pavich, M., and Caffee, M. (accepted) Rates of sediment supply to arroyos from uplands determined using in situ produced cosmogenic ^{10}Be and ^{26}Al in sediments. **Quaternary Research**.
- Nichols, K.K., and Bierman, P.R. (accepted) Fifty four years of ephemeral channel response to intense military activity at Camp Iron Mountain, Mojave Desert, California, Environmental and Engineering Impacts of Military Operations, **Geological Society of America Reviews of Engineering Geology**.
- Colgan, P.M., Mickelson, D. M., Bierman, P. R., Caffee, M. W., (in review) Variation in glacial erosion near the southern margin of the Laurentide Ice Sheet, south central Wisconsin, USA: implications for cosmogenic dating of glacial terrains: **Geological Society of America Bulletin**
- Schroeder, P.A., Nathan D., Melear, N.D., Bierman, P.R., Kashgarian, M., and Caffee, M.W., (accepted) Apparent gibbsite growth ages for the regolith in the Georgia Piedmont, **Geochimica et Cosmochimica Acta**.
- Bierman, P. R., Marsella, K. A., Davis, P. T., Patterson, C. and Caffee, M., (1999), Mid-Pleistocene cosmogenic minimum-age limits for pre-Wisconsinan glacial surfaces in southwestern Minnesota and southern Baffin Island -- a multiple nuclide approach. **Geomorphology**, 27, n 1 / 2, p. 25-40.

Book chapters

- Bierman, P. (in press) In situ produced cosmogenic isotopes: dating landforms and monitoring erosion rates. **Encyclopedia of Quaternary Science**.
- Bierman, P.R., Albrecht, A., Bothner, M., Brown, E., Bullen, T., Gray, L., Turpin, L., (1998) Weathering, erosion and sedimentation, In: Kendall, C., and McDonnell, J. J., **Isotope Tracers in Catchment Hydrology**, chapter 23, Elsevier

Conference Proceedings

- Clapp, E.M., Bierman, P.R., Caffee, M.W. (1999) Sediment generation and export rates in the Nahal Yale drainage basin, determined from cosmogenic ^{10}Be and ^{26}Al , Negev Desert, southern Israel, in **Drainage Basin dynamics and morphology, conference excursion, Negev Desert**, Lekach, J. and Hassan, M.A, eds., Hebrew University Jerusalem, p. 98-103.

Abstracts

- Bierman, P., Clapp, E., Caffee, M., and Schroeder, P. (1999) Understanding Earth surface processes with 10-Be (and a little 26-Al), **Geological Society of America Abstracts with Programs**. (National).
- Bierman, P., Clapp, E., Massey, C., and, Caffee, M., (1999) Tracing sediment through drainage basins with cosmogenic radionuclides, ^{10}Be and ^{26}Al . **International Conference on Drainage basin Dynamics and Morphology**, Jerusalem (International)
- Clapp, E., Bierman, P.R., and Caffee, M., (1999), Sediment generation and export rates in the Nahal Yael drainage basin, determined from cosmogenic ^{10}Be and ^{26}Al , Negev desert, southern Israel, **Geological Society of America Abstracts with Programs**, 31 (7), A-256. (National)
- Nichols, K.K, Bierman, P.R., Caffee, M. (1999) Long-term sediment dynamics of the Iron and Granite Mountain piedmonts, Mojave Desert, California, U.S.A., **International Conference on Drainage basin Dynamics and Morphology**, Jerusalem, Israel, p. 47
- Nichols, K. K. and Bierman, P.R. (1998). Geomorphic response to military training in the Mojave Desert. **Geological Society of America Abstracts with Programs**, 30 (7), A-143 (National)
- Clapp, E., Bierman, P. and, Caffee, M., (1998) Estimating long-term erosion rates in a hyper-arid region using in situ produced cosmogenic 10-Be and 26-Al in sediment and bedrock. **Geological Society of America Abstracts with Programs**, 30 (7), A-361. (National)
- Clapp, E. M. and Bierman, P. R. (1997) Rates of erosion determined using in situ produced cosmogenic isotopes in a small arroyo basin, northwestern New Mexico **Geological Society of America Abstracts with Programs**, 29 (7), A-371 (National).

Advanced degrees Earned by Project Personnel

Kyle Nichols, MS, 2000

Erik Clapp, Ph.D., ABD (expected completion, Fall 2000)

Students and faculty supported in whole or in part by ARO funding

Paul Bierman, Professor, Geology Department, University of Vermont, UVM

Erik Clapp, Doctoral Candidate, School of Natural Resources, UVM, ABD

Lyman Persico, BS (Honors) , Geology, UVM, Candidate

Kyle Nichols, MS, Geology, UVM, 1999

Darrin Santos, BS (Honors), Environmental Science, UVM, 1999

Sara Gran, MS. Geology, UVM, 1999

Sean Lampert, BS, Geology, UVM, 1997

Ben Copans, BS, Geology, UVM, 1998